



Tailoring ZSM-5 zeolites for the fast pyrolysis of biomass to aromatic hydrocarbons

Hoff, Thomas C.; Gardner, David W.; Thilakaratne, Rajeeva; Wang, Kaige; Hansen, Thomas W.; Brown, Robert C.; Tessonnier, Jean-Philippe

Publication date:
2016

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
Hoff, T. C., Gardner, D. W., Thilakaratne, R., Wang, K., Hansen, T. W., Brown, R. C., & Tessonnier, J-P. (2016). *Tailoring ZSM-5 zeolites for the fast pyrolysis of biomass to aromatic hydrocarbons*. Abstract from 252nd ACS National Meeting, Philadelphia, Pennsylvania, United States.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

CATL - Catalysts & Catalytic Technologies for Conversion of Biomass & Its Derivatives

Tailoring ZSM-5 zeolites for the fast pyrolysis of biomass to aromatic hydrocarbons

Thomas C. Hoff, David W. Gardner, Rajeeva Thilakaratne, Kaige Wang, Thomas W. Hansen, Robert C. Brown, Jean-Philippe Tessonier

1. Chemical and Biological Engineering, Iowa State University, Ames, IA, United States.
2. Bioeconomy Institute, Iowa State University, Ames, IA, United States.
3. Center for Electron Nanoscopy, Technical University of Denmark, Lyngby, Denmark

Catalytic fast pyrolysis (CFP) represents a sustainable method to convert raw biomass from different sources into useful chemicals and fuels in a single step. ZSM-5 zeolite has been identified as the best catalyst for this reaction. Its pore structure provides unique shape selectivity toward C₆-C₁₁ aromatics—in particular to benzene, toluene, xylenes—which are important building blocks for the chemical industry. However, the exact role and impact of catalyst features such as crystallinity, elemental composition, porosity, and acidity on the catalyst's performance remain elusive. The interplay of these parameters under reaction conditions represents a major roadblock that has hampered significant improvement in catalyst design for over a decade.

Here, we studied commercial and laboratory synthesized ZSM-5 zeolites and combined data from ten complementary characterization techniques in an attempt to identify parameters common to high-performance catalysts. We show that crystallinity and framework aluminum sites accessibility are critical to achieve high aromatic yields. In contrast, mesoporosity which was expected to enhance intra-crystalline diffusion played only a minor role. These findings enabled us to synthesize a ZSM-5 catalyst with enhanced activity, offering the highest aromatic hydrocarbon yield reported to date.